

---

Preoperative:

Propel treatment accelerates the speed of your orthodontic treatment by 50%.

Prior to your treatment, please note the following:

- Do **NOT** take Advil or Motrin 24 hours before treatment
- You will be numb for the treatment, please plan accordingly

Postoperative:

After treatment, it is normal to experience any of the following:

- Minor bleeding and redness at the treatment sites
- Tenderness for 24-28 hours after treatment
- Soreness with teeth during aligner changes

Take Tylenol for discomfort. Do **NOT** take Advil or Motrin.

Unless otherwise instructed, continue to brush and floss as usual

Instructions for Invisalign Trays:

- Wear aligners for 22-23 hours a day for \_\_\_\_\_ days as directed and follow aligner schedule as provided
- Call our office if your aligners do not fit the same as they do during initial visit
- Call our office if you lose or break an aligner
- Call our office if you notice a lost button on any tooth/teeth
- Take aligners out to eat and brush teeth prior to placing aligners back in
- Take aligners out to drink any **HOT** or not clear liquids
- Make sure you keep aligners in your mouth or in their case. Do not leave aligners anywhere else

Additional treatment may be necessary throughout your Invisalign treatment and that will be determined with an intraoral progress scan and deemed necessary by Dr. DiSano.

---

If you have questions or concerns, please call the office so we may address them on a case-by-case basis.

**401 783-1530**